

Analyticity, Crossing Symmetry and the Limits of Chiral Perturbation Theory

R. Sekhar Chivukula^{a,1}

Michael J. Dugan^{a,b,2}

and

Mitchell Golden^{a,3}

The chiral Lagrangian for Goldstone boson scattering is a power series expansion in numbers of derivatives. Each successive term is suppressed by powers of a scale, Λ_χ , which must be less than of order $4\pi f/\sqrt{N}$ where f is the Goldstone boson decay constant and N is the number of flavors. The chiral expansion therefore breaks down at or below $4\pi f/\sqrt{N}$. We argue that the breakdown of the chiral expansion is associated with the appearance of physical states other than Goldstone bosons. Because of crossing symmetry, some “isospin” channels will deviate from their low energy behavior well before they approach the scale at which their low energy amplitudes would violate unitarity. We argue that the estimates of “oblique” corrections from technicolor obtained by scaling from QCD are untrustworthy.

6/92

^aBoston University, Department of Physics, 590 Commonwealth Avenue, Boston, MA 02215

^bLyman Laboratory of Physics, Harvard University, Cambridge, MA 02138

¹SEKHAR@WEYL.BU.EDU

²DUGAN@HUHEPL.BITNET

³GOLDEN@WEYL.BU.EDU

1. Introduction

The chiral Lagrangian [1] is a compact way of calculating the amplitudes for low-energy processes involving Goldstone bosons. Though it was originally developed to describe pions, it has more recently been applied to calculate amplitudes for processes involving the longitudinal components of the W and Z gauge bosons [2] [3] [4]. The equivalence theorem [3] insures that, at energies large compared to M_W , the scattering amplitudes of longitudinally polarized gauge bosons are approximately the same as those of the Goldstone bosons which would be present in the ungauged theory. In technicolor models [5], frequently there are also additional, approximate Goldstone bosons [6]. For example, the one-family model [7] has 60 pseudo-Goldstone bosons, in addition to the 3 “swallowed” degrees of freedom. In this case, the chiral Lagrangian may also be used to describe low-energy processes involving pseudo-Goldstone bosons as well as longitudinal W or Z bosons [8].

Consider a model in which the symmetry breaking pattern is $SU(N)_L \times SU(N)_R \rightarrow SU(N)_V$ ($N = 2$ in the simplest technicolor model and $N = 8$ in the one-family model). The most general chirally invariant Lagrangian [1] may be written in terms of the field

$$\Sigma = \exp(2i\pi^a T^a / f) , \quad (1.1)$$

where the π^a (which we refer to as “pions”) are the Goldstone boson fields, the T^a are the generators of $SU(N)$, normalized to $\text{tr} T^a T^b = \delta^{ab}/2$, and f is the analog of the pion decay constant. Under a chiral transformation, the field Σ transforms as $\Sigma \rightarrow L\Sigma R^\dagger$, with $L \in SU(N)_L$ and $R \in SU(N)_R$. The most general chirally invariant Lagrangian can be written as an expansion in powers of derivatives. There are no nontrivial chirally invariant terms involving no derivatives and only

$$\mathcal{L}_2^{(0)} = \frac{f^2}{4} \text{tr} \partial^\mu \Sigma^\dagger \partial_\mu \Sigma \quad (1.2)$$

with two derivatives. Additional terms with more derivatives are suppressed by powers of some momentum scale, denoted Λ_χ .

Chiral perturbation theory is an expansion in p^2/Λ_χ^2 . The utility of the chiral Lagrangian arises from the fact that at energies less than Λ_χ , the interactions of exact Goldstone bosons are determined by $\mathcal{L}_2^{(0)}$, *i.e.* they are entirely determined by the symmetry structure of the theory. For this reason, these lowest order predictions are universal [4].

At energies near or above Λ_χ , however, all terms in the expansion contribute and the chiral Lagrangian becomes effectively useless. Here the amplitudes become model

dependent. The expansion must fail at or before an energy scale associated with the appearance of additional particles or resonances: for example, in the case of the one Higgs-doublet standard model the chiral expansion fails at an energy scale of order the Higgs mass. In a weakly coupled theory, these particles will generally be light and the chiral expansion will fail at low energies.

In a strongly interacting theory, such as QCD or technicolor, the resonances will generally be heavy, and the predictions of chiral perturbation theory may be used to fairly high energies. It is important, therefore, to understand how large Λ_χ may be. This question was first addressed by Weinberg [9] who argued that, since the higher order terms are required as counterterms to loops involving the lowest order interactions, it was inconsistent to assume that the size of these terms was smaller than that typical of the corresponding loop correction. This leads to the estimate of naive dimensional analysis that Λ_χ must be less than or about $4\pi f$.

In this paper, we discuss how the bound on Λ_χ varies with N , *i.e.* how it depends on the number of Goldstone bosons. By examining the size of corrections to the one-loop effective action, Sundrum and Soldate [10] have argued that Λ_χ is bounded by $4\pi f/\sqrt{N}$. We address this question by examining pion scattering in chiral perturbation theory, and explicitly confirm their results. This result does not depend on any expansion in $1/N$. In particular, if N is significantly larger than 2 (*e.g.* 8 in the one-family model), the chiral Lagrangian ceases to be valid at energies much lower than suggested by naive dimensional analysis. This change is particularly important in the context of electroweak symmetry breaking phenomenology at the SSC and LHC [11] [8][12].

As previously stated, the chiral symmetries of technicolor are generally only approximate symmetries. In addition to the three “eaten”, exact Goldstone bosons, there are often additional pseudo-Goldstone bosons. If the mass of a typical pseudo-Goldstone boson is m , chiral perturbation theory is also an expansion in m^2/Λ_χ^2 , *i.e.* Λ_χ also sets the scale for the size of corrections due to chiral symmetry breaking. Chiral symmetry breaking interactions induce nonderivative terms in the chiral Lagrangian. For simplicity, in this paper we consider a chiral symmetry breaking interaction which gives the same mass to all Goldstone bosons. At lowest order in the symmetry breaking the Lagrangian is

$$\mathcal{L}_2 = \mathcal{L}_2^{(0)} + \frac{m^2 f^2}{4} \text{tr}(\Sigma + \Sigma^\dagger) . \quad (1.3)$$

We will show that in a theory with a large number of pseudo-Goldstone bosons, since Λ_χ is smaller than expected, the effects of chiral symmetry breaking are correspondingly

larger than expected. While this particular kind of symmetry breaking cannot occur in a technicolor theory – such a symmetry breaking term would break the weak gauge symmetry – the calculation will illustrate the enhancement of symmetry breaking effects in a theory with $N > 2$.

In the next section we consider Goldstone boson scattering at tree level using $\mathcal{L}_2$. Using crossing invariance we show that all amplitudes may be expressed in terms of two invariant functions. Decomposing the amplitude into flavor and angular momentum channels, we find that the $SU(N)_V$ singlet, spin zero channel would violate unitarity at an energy of order $4\pi f/\sqrt{N}$. As physical amplitudes cannot violate unitarity, we can conclude that that terms of order p^4 and higher must become important and that Λ_χ cannot be larger than of order $4\pi f/\sqrt{N}$. This argument is a generalization of that applied by Lee, Quigg, and Thacker [13] to longitudinal gauge boson scattering in the standard one Higgs-doublet model.

In the third section, we compute the corrections to Goldstone boson scattering at one-loop. Confirming ref. [10], we find that the corrections are larger than naively expected by a factor of N . We give a simple argument that this pattern persists to all orders in the loop expansion implying again that Λ_χ is of order $4\pi f/\sqrt{N}$. This implies that *all* terms in the chiral expansion are relevant at these energies and chiral perturbation theory breaks down.

While the singlet channel saturates unitarity at energies of order $4\pi f/\sqrt{N}$, the other channels have amplitudes which are still much less than one. Is it possible that the low energy predictions from $\mathcal{L}_2$ continue to be valid in these other channels, as in a generalization of the “conservative” model of Chanowitz and Gaillard in [3]? In the fourth section, we argue that in general, the chiral Lagrangian will be valid at all energies below the energy scale associated with new physical states, *i.e.* Λ_χ may always be interpreted as the scale of new physics and is not just a formal artifact of chiral perturbation theory. This new physics will enter in all channels and hence the low-energy predictions for *all* channels fail. The behavior of the theory above Λ_χ depends on what new physics is present, and one cannot trust arbitrary unitarizations of the low-energy amplitudes (such as summing a subset of the chiral loop diagrams, the K -matrix, or the Padé approximants) [14][15].

In QCD, the chiral Lagrangian ceases to be valid at an energy scale of order the mass of the ρ meson. The arguments we make here imply that in a theory with more than two light flavors, at least some of the resonances are lighter than would be expected by scaling from QCD, and that any results for such a theory based on scaling from QCD are suspect. In particular, the estimates of “oblique” corrections in the one-family technicolor model [16] are not trustworthy.

2. Pion Scattering from $\mathcal{L}_2$

Consider the scattering process $\pi^a \pi^b \rightarrow \pi^c \pi^d$. The amplitude for such a process may be decomposed into irreducible $SU(N)_V$ representations. We may understand the representations contained in **adjoint** $\otimes$ **adjoint** of $SU(N)$ as follows. Consider the object $(T^a)_j^i (T^b)_l^k$, where the indices a, b are in the **adjoint** of $SU(N)$ while i, j, k, l are in the fundamental. First, we may trace i with l , and j with k , yielding a singlet. Next, we may form an **adjoint**: remove the singlet already constructed, then symmetrize (i, k) , antisymmetrize (j, l) , and contract i with l . Another **adjoint** is obtained by symmetrizing both (i, k) and (j, l) before contracting. The next possibility is to antisymmetrize (i, k) and (j, l) , and remove all traces. This representation has dimension $(N+1)(N-3)N^2/4$. We may symmetrize (i, k) and (j, l) and remove traces, yielding a representation of dimension $(N-1)(N+3)N^2/4$. Lastly there are two complex representations, conjugates of each other, in which the traces are removed and either (i, k) is symmetric and (j, l) is antisymmetric or vice-versa. These complex representations have dimension $(N+2)(N+1)(N-1)(N-2)/4$. We refer to the seven representations as Δ , F , D , Y , X , T , and $\bar{T}$, respectively. The representations Δ , D , Y , and X are symmetric under the exchange of a and b , and therefore only the angular momentum $J = \text{even}$ partial waves can contribute. The others are antisymmetric under $a \leftrightarrow b$, and only the odd angular momentum partial waves contribute. For $N = 2$ only Δ , F , and X exist, corresponding to the isospin 0, 1, and 2 channels. If $N = 3$, Y does not exist, while the others are respectively the **1**, **8_a**, **8_s**, **27**, **10**, and **10**.

Next, we construct the most general amplitude for $\pi\pi$ scattering consistent with Bose symmetry, crossing invariance, and $SU(N)_V$ conservation. There are nine invariant tensors with four adjoint indices, corresponding to the nine singlets in **(adjoint)**⁴. A set of nine¹ linearly independent invariants is $\delta^{ab}\delta^{cd}$, $\delta^{ac}\delta^{bd}$, $\delta^{ad}\delta^{bc}$, $d^{abe}d^{cde}$, $d^{ace}d^{bde}$, $d^{ade}d^{bce}$, $d^{abe}f^{cde}$, $d^{ace}f^{bde}$, and $d^{ade}f^{bce}$, where d^{abc} and f^{abc} are defined by

$$f^{abc} = -2i\text{tr}[T^a, T^b]T^c \quad \text{and} \quad d^{abc} = 2\text{tr}\{T^a, T^b\}T^c. \quad (2.1)$$

There can be no part of the amplitude proportional to $d^{abe}f^{cde}$: because of Bose symmetry and the symmetry of d^{abc} under $a \leftrightarrow b$, the initial state would have to be in an even angular

¹ There are nine when $N > 3$. Using the relations in Appendix A, all the other invariants can be shown to dependent on these nine. For $N = 2$ the d symbols do not exist and there are only three invariants. For $N = 3$ the relation (Burgoyne's identity [17]) $3(d^{abe}d^{cde} + d^{ace}d^{bde} + d^{ade}d^{bce}) = \delta^{ab}\delta^{cd} + \delta^{ac}\delta^{bd} + \delta^{ad}\delta^{bc}$ permits us to eliminate one of the invariants.

momentum state, while by the antisymmetry of f^{cde} , the final state would have to be in an odd angular momentum state. Therefore, the most general amplitude is

$$a(s, t, u)^{a,b;c,d} = \delta^{ab}\delta^{cd}A(s, t, u) + \delta^{ac}\delta^{bd}A(t, s, u) + \delta^{ad}\delta^{bc}A(u, t, s) \\ + d^{abe}d^{cde}B(s, t, u) + d^{ace}d^{bde}B(t, s, u) + d^{ade}d^{bce}B(u, t, s) , \quad (2.2)$$

where s , t , and u are the Mandelstam variables and A and B are unknown functions. Bose symmetry also implies that the functions A and B must be symmetric under the exchange of their second and third arguments.

Applying the projection operators defined in Appendix A to the amplitude (2.2), the amplitudes for pion scattering in the various $SU(N)_V$ channels can all be written in terms of A and B :

$$\begin{aligned} a_\Delta(s, t, u) &= (N^2 - 1)A(s, t, u) + A(t, s, u) + A(u, t, s) \\ &\quad + \frac{(N^2 - 4)}{N}(B(t, s, u) + B(u, t, s)) \\ a_F(s, t, u) &= A(t, s, u) - A(u, t, s) + \frac{N^2 - 4}{2N}(B(t, s, u) - B(u, t, s)) \\ a_D(s, t, u) &= A(t, s, u) + A(u, t, s) + \frac{N^2 - 4}{N}B(s, t, u) \\ &\quad + \frac{N^2 - 12}{2N}(B(t, s, u) + B(u, t, s)) \\ a_Y(s, t, u) &= A(t, s, u) + A(u, t, s) - \frac{N + 2}{N}(B(t, s, u) + B(u, t, s)) \\ a_X(s, t, u) &= A(t, s, u) + A(u, t, s) + \frac{N - 2}{N}(B(t, s, u) + B(u, t, s)) \\ a_T(s, t, u) &= a_{\bar{T}}(s, t, u) = A(t, s, u) - A(u, t, s) - \frac{2}{N}(B(t, s, u) - B(u, t, s)) \end{aligned} \quad (2.3)$$

The lowest order chiral Lagrangian gives $A(s, t, u) = (2/N)(s - m^2)/f^2$, and $B(s, t, u) = (s - m^2)/f^2$.

The partial wave amplitudes are defined by

$$a_{I\ell}(s) = \frac{1}{64\pi} \int_{-1}^1 a_I(s, \cos \theta) P_\ell(\cos \theta) d \cos \theta , \quad (2.4)$$

where P_ℓ is the Legendre polynomial of order ℓ . The scattering amplitudes in the various

channels from $\mathcal{L}_2$ [14] are then:

$$\begin{aligned}
a_{\Delta 0} &= \frac{Ns}{32\pi f^2} - \frac{m^2}{16\pi N f^2} \\
a_{F1} &= \frac{Ns}{192\pi f^2} - \frac{Nm^2}{48\pi f^2} \\
a_{D0} &= \frac{Ns}{64\pi f^2} - \frac{m^2}{8\pi N} \\
a_{Y0} &= \frac{s}{32\pi f^2} - \frac{m^2}{16\pi f^2} \\
a_{X0} &= -\frac{s}{32\pi f^2} + \frac{m^2}{16\pi f^2} .
\end{aligned} \tag{2.5}$$

All other partial wave amplitudes are zero (including a_{T1} and $a_{\bar{T}1}$). Note that $a_{\Delta 0}$, a_{F1} and a_{D0} are enhanced by a factor of N .

The amplitude $a_{\Delta 0}$ calculated at tree level is real, and (for small m^2) would exceed 1 when $\sqrt{s} > 4\pi f/\sqrt{N}$. A physical scattering amplitude must lie on or inside the Argand circle. At these energies, therefore, loop corrections and higher order terms in the chiral Lagrangian must make as large a contribution as the two-derivative term, and the calculation using $\mathcal{L}_2$ ceases to be useful. This suggests that Λ_χ is less than or of order $4\pi f/\sqrt{N}$, as was emphasized in [10]. Note that this result holds independent of the largeness of N . No expansion in powers of $1/N$ need be made.

3. Pion Scattering at Order p^4

An alternative approach to put a limit Λ_χ is based on an estimate of the size of loop corrections [9]. Since the theory is not renormalizable, the terms of order p^4 are required as counterterms to loops involving the lowest order interactions. In calculating the scattering amplitude to order p^4 , one must consider tree-level diagrams with interactions coming from operators of fourth order in momenta, and one-loop diagrams using $\mathcal{L}_2$. It is unnatural to assume that the contribution from the former is much larger than the latter since such a statement could only be true for a particular choice of renormalization scale. Similarly, the two-loop calculation using $\mathcal{L}_2$ will require counterterms of order p^6 , etc. In this section we compute the one-loop corrections to Goldstone boson scattering.

The next-to-leading order Lagrangian is made up of terms containing four derivatives, two derivatives and one power of the symmetry breaking parameter, or two powers of the symmetry breaking:

$$\begin{aligned}
\mathcal{L}_4 = & \ell_1 \{ \text{tr}(\partial^\mu \Sigma^\dagger \partial_\mu \Sigma) \}^2 + \ell_2 \{ \text{tr}(\partial^\mu \Sigma^\dagger \partial^\nu \Sigma) \}^2 + \\
& \ell_3 \text{tr}(\partial^\mu \Sigma^\dagger \partial_\mu \Sigma \partial^\nu \Sigma^\dagger \partial_\nu \Sigma) + \ell'_3 \text{tr}(\partial^\mu \Sigma^\dagger \partial^\nu \Sigma \partial_\mu \Sigma^\dagger \partial_\nu \Sigma) + \\
& \ell_4 m^2 \text{tr}(\Sigma + \Sigma^\dagger) \text{tr}(\partial^\mu \Sigma^\dagger \partial_\mu \Sigma) + \ell_5 m^2 \text{tr}\{(\Sigma + \Sigma^\dagger)(\partial^\mu \Sigma^\dagger \partial_\mu \Sigma)\} + \\
& \ell_6 m^4 \{ \text{tr}(\Sigma + \Sigma^\dagger) \}^2 + \ell_7 m^4 \{ \text{tr}(\Sigma - \Sigma^\dagger) \}^2 + \ell_8 m^4 \text{tr}(\Sigma^2 + \Sigma^{\dagger 2}) .
\end{aligned} \tag{3.1}$$

All other possible terms vanish by the equations of motion. This notation agrees with that of Gasser and Leutwyler[18]. In their case the term proportional to ℓ'_3 is not linearly independent, because they were considering an $SU(3) \times SU(3)$ chiral symmetry.

The calculation of the $\pi\pi \rightarrow \pi\pi$ amplitudes at one loop is straightforward, if tedious.

The result is:

$$\begin{aligned}
A(s, t, u) = & \frac{2}{N} \frac{s - m^2}{f^2} \\
& + \frac{J(s)}{f^4} \left(-\frac{1}{2} s^2 + \frac{2}{N^2} m^4 \right) + \frac{J(t)}{f^4} \left(-\frac{1}{3} t^2 + \frac{5}{3} m^2 t - \frac{7}{3} m^4 - \frac{1}{6} st + \frac{2}{3} m^2 s \right) \\
& + \frac{J(u)}{f^4} \left(-\frac{1}{3} u^2 + \frac{5}{3} m^2 u - \frac{7}{3} m^4 - \frac{1}{6} su + \frac{2}{3} m^2 s \right) \\
& + \frac{s^2}{f^4} \left(-\frac{2}{3} \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} + 8\ell_1(\mu) + 4\ell_2(\mu) + \frac{8}{N} \ell_3(\mu) + \frac{23}{18} \frac{1}{16\pi^2} \right) \\
& + \frac{tu}{f^4} \left(\frac{2}{3} \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} - 8\ell_2(\mu) - \frac{16}{N} \ell'_3(\mu) - \frac{13}{9} \frac{1}{16\pi^2} \right) \\
& + \frac{m^2 s}{f^4} \left(\frac{2}{3} \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} - 32\ell_1(\mu) - 16\ell_2(\mu) - \frac{32}{N} \ell_3(\mu) \right. \\
& \quad \left. + 16\ell_4(\mu) + \frac{16}{N} \ell_5(\mu) - \frac{28}{9} \frac{1}{16\pi^2} \right) \\
& + \frac{m^4}{f^4} \left(\left(-2 + \frac{2}{N^2} \right) \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} + 32\ell_1(\mu) + 32\ell_2(\mu) + \frac{32}{N} \ell_3(\mu) + \frac{32}{N} \ell'_3(\mu) \right. \\
& \quad \left. - 32\ell_4(\mu) - \frac{32}{N} \ell_5(\mu) + 32\ell_6(\mu) + \frac{32}{N} \ell_8(\mu) + \left(\frac{20}{3} - \frac{4}{N^2} \right) \frac{1}{16\pi^2} \right)
\end{aligned} \tag{3.2}$$

$$\begin{aligned}
B(s, t, u) = & \frac{s - m^2}{f^2} \\
& + \frac{NJ(s)}{f^4} \left(-\frac{1}{8}s^2 + \frac{2}{N^2}m^4 \right) + \frac{NJ(t)}{f^4} \left(-\frac{1}{24}t^2 + \frac{1}{3}m^2t - \frac{2}{3}m^4 - \frac{1}{12}st + \frac{1}{3}m^2s \right) \\
& + \frac{NJ(u)}{f^4} \left(-\frac{1}{24}u^2 + \frac{1}{3}m^2u - \frac{2}{3}m^4 - \frac{1}{12}su + \frac{1}{3}m^2s \right) \\
& + \frac{Ns^2}{f^4} \left(-\frac{1}{12}\frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} + \frac{4}{N}\ell_1(\mu) + \frac{5}{36}\frac{1}{16\pi^2} \right) \\
& + \frac{Ntu}{f^4} \left(+\frac{1}{12}\frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} - \frac{8}{N}\ell'_3(\mu) - \frac{2}{9}\frac{1}{16\pi^2} \right) \\
& + \frac{Nm^2s}{f^4} \left(-\frac{1}{6}\frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} - \frac{16}{N}\ell_3(\mu) + \frac{8}{N}\ell_5(\mu) - \frac{5}{9}\frac{1}{16\pi^2} \right) \\
& + \frac{Nm^4}{f^4} \left(\frac{2}{N^2}\frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} + \frac{16}{N}\ell_3(\mu) + \frac{16}{N}\ell'_3(\mu) - \frac{16}{N}\ell_5(\mu) + \frac{16}{N}\ell_8(\mu) \right. \\
& \quad \left. + \left(\frac{4}{3} - \frac{4}{N^2} \right) \frac{1}{16\pi^2} \right)
\end{aligned} \tag{3.3}$$

Here we have used dimensional regularization in $4 - \epsilon$ dimensions with a scale μ and, as detailed in Appendix B, the parameters $\ell_i(\mu)$ are renormalized at this scale. The function J is defined as

$$J(x) = \frac{1}{16\pi^2} \sqrt{1 - \frac{4m^2}{x}} \log \frac{\sqrt{1 - \frac{4m^2}{x}} + 1}{\sqrt{1 - \frac{4m^2}{x}} - 1} \tag{3.4}$$

where $\log(z)$ and $\sqrt{z}$ are both understood to have a cut under the negative real axis. The quantities m and f which appear in (3.2) and (3.3) are the physical mass and decay constant. Their renormalizations are also given in Appendix B.

From (3.2) and (3.3) we see that the typical corrections to the tree level results are of order $Ns/(16\pi^2 f^2)$ or $Nm^2/(16\pi^2 f^2)$. In agreement with the previous section and ref. [10], then, it is inconsistent to assume that Λ_χ is much greater than $4\pi f/\sqrt{N}$.

This pattern, *i.e.* that loop results are larger by powers of N than expected by naive dimensional analysis, persists to all orders. In order to keep track of N , we may use a double line notation for each Goldstone boson, analogous to the double line notation used by 't Hooft [19] in large- N_C QCD. Each Goldstone boson is a member of the adjoint representation of $SU(N)_V$, and is analogous to a gluon in the adjoint of the $SU(N_C)$ color. Analogous to large- N_C QCD, then, the contributions to a process at L loops using

interactions from $\mathcal{L}_2$ that have the highest power of N are proportional to $N^L/(16\pi^2)^L$. Therefore, there are L -loop corrections to pion scattering of order

$$\left(\frac{\sqrt{N}p}{4\pi f}\right)^{2L}, \quad (3.5)$$

where p is a typical momentum in the process. Since all higher momentum corrections are suppressed by the same energy scale, chiral perturbation theory as a whole breaks down by energies of order $4\pi f/\sqrt{N}$. For any fixed N , it is *possible* that the subleading in N diagrams are numerically as important as those that are leading. However, making Λ_χ much bigger than $4\pi f/\sqrt{N}$ would require both an enhancement at every loop of the subleading diagrams by a numerical factor as large as N , and a cancellation of these with the leading diagrams.

4. Implications of Analyticity and Crossing

Some interesting questions arise at this point. We have argued that chiral perturbation theory breaks down at or before Λ_χ , but what actually happens to the amplitudes as s increases beyond this value? What is the significance of Λ_χ ? The amplitudes for the partial waves other than $a_{\Delta 0}$ are all below their unitarity limits when $\sqrt{s} = 4\pi f/\sqrt{N}$. Is it possible that these other channels continue to behave like the prediction of the lowest order chiral Lagrangian, as in the $SU(N)_V$ generalization of the “conservative model” of ref. [3]?

The pion scattering amplitudes are determined by the two functions A and B . We will examine the analytic structure of these functions in the complex s, t hypersurface, where u is determined by the mass shell condition. Let us consider the case of a small $m^2 > 0$, so as to avoid the subtleties associated with infrared problems. Amplitudes in chiral perturbation theory are expansions in s and t . The S -matrix is analytic on this hypersurface except for cuts on the physical (s, t) plane and poles or cuts on the unphysical hypersheets. The cuts in the physical (s, t) plane are due to multipion states, and the appearance of poles or other structure on the unphysical hypersheets correspond to resonances or physical states other than pions. The masses of the new physical states are not protected by a chiral symmetry and the corresponding structures in the S -matrix are therefore away from the origin.

Consider the region R with s , t , and u all less than $4m^2$. Here, assuming there are no particles lighter than twice the pion mass, the S -matrix is analytic. The functions A and B computed from the chiral Lagrangian to arbitrary order have an infinite set of adjustable coefficients multiplying the terms $s^i t^j$ for all i and j . We can regard this as a convergent expansion about any point in the region R . Therefore, the amplitude computed in chiral perturbation theory can be adjusted to match the S -matrix exactly in R . Going outside R , the singularities of the S -matrix on the physical plane correspond to cuts from multipion states and are determined by unitarity. They are correctly included in chiral loop calculations. Moving away from this region, therefore, the chiral Lagrangian calculation should reproduce the S -matrix so long as there aren't any singularities closer to the origin associated with physical states *other* than multipion states. That is, chiral perturbation theory is a good approximation to the S -matrix *all the way* out to an energy scale associated with the appearance of new physics.

Admittedly, the argument given above for the range of validity of the chiral expansion is not truly rigorous. This is because it works only if one computes without expanding in m^2 . As emphasized by Pagels and Li [20], the S -matrix is not analytic in m^2 . However, as motivated by the theory of critical phenomena [21], we assume that the coefficients of the Chiral Lagrangian are analytic in m^2 and that the nonanalyticity of the S -matrix is reproduced by chiral loop calculations.

If this standard assumption is correct, then the arguments given above are correct so long as m^2 is small enough, and the chiral expansion is valid for all energies below the first nonanalytic structure in the S -matrix corresponding to some real new physics. The breakdown of the chiral Lagrangian is not a calculational artifact; the scale Λ_χ has direct physical significance, unlike the scale Λ_{QCD} in perturbation theory. Moreover, the arguments above show that this physical scale cannot be larger than about $4\pi f/\sqrt{N}$.

In general, there are many possibilities for the new physics at Λ_χ . In QCD with $N = 2$, the ρ pole gives a singularity in A when t or u is $m_\rho^2 - im_\rho\Gamma_\rho$. In the one-doublet Higgs model, there is a pole in A whenever s is $M_H^2 - iM_H\Gamma_H$. The first singularity to appear could be a branch cut instead of a pole. Consider a world in which the π s are massless, but the K s weigh 200 MeV. (The pion decay constant is still 93 MeV.) Imagine constructing an $SU(2)$ invariant chiral Lagrangian to describe the massless pions. The first new singular structure one encounters in the S -matrix is the two K branch cut, at 400 MeV, and at this energy the chiral expansion breaks down.

Since the scale Λ_χ is associated with non-universal structure, such as the ρ , the Higgs, or the KK states in the examples above, the behavior of the scattering amplitudes at energies at or above Λ_χ *cannot be* universal. Therefore, there is no reason to trust predictions based on an arbitrary, *e.g.* K -matrix, Padé approximant[15], or bubble sum[14], unitarization of the universal, lowest-order, chiral amplitudes.

Finally, we address the issue of whether the channels other than $a_{\Delta 0}$ can follow their low-energy predictions beyond energies of order Λ_χ . Consider again the functions A and B . Because of the crossing relations (2.3), all channels will be affected by whatever new physics enters at the scale Λ_χ . In particular we expect that at Λ_χ the amplitudes in all channels will deviate strongly from the low-energy predictions, even though the low-energy predictions may be well below 1.

Consider pion scattering in QCD. The data² and lowest-order predictions for the low isospin channels are plotted in figs. 1,2,3. The amplitude in the isospin-0 spin-0 channel ($a_{\Delta 0}$) starts to deviate from the lowest order prediction at an energy of about 600 MeV. This is not surprising, because the amplitude is at that point a substantial fraction of its unitarity limit. Examining the other partial waves, we find that they deviate from the lowest order chiral Lagrangian formulas above approximately *the same energy scale*. For example, in the isospin-1 spin-1 channel (a_{F1}), the ρ resonance appears at 770 MeV. While we cannot predict the appearance of the ρ , the fact that the isospin 1 and 2 amplitudes deviate strongly from their low-energy predictions while they would still be relatively weakly coupled is not a surprise.

5. Conclusions and Speculations

In this paper, we have argued that in general, the chiral expansion will be valid at all energies below the energy scale associated with new physical states. That is, Λ_χ may always be interpreted as the scale of new physics and is not just a formal artifact of chiral perturbation theory. In general, there is no procedure to infer the high-energy structure of the theory from the universal lowest-order chiral lagrangian. In the three examples given in section four, the high-energy physics is different, while the low-energy behavior is precisely the same. In particular, there is no reason to trust the amplitudes constructed by unitarizing the lowest-order predictions.

² We thank G. Valencia for providing us with the data compiled for reference [22].

In a theory with a spontaneously broken $SU(N)_L \times SU(N)_R$ chiral symmetry, the scale Λ_χ is bounded by $4\pi f/\sqrt{N}$. This result depends only on the low-energy effective theory and is independent of the precise form of the fundamental theory. If the theory is QCD-like, Λ_χ is presumably associated with the appearance of resonances like the ρ . When $N = 2$ and $f_\pi = 93$ MeV, the limit on Λ_χ is about 825 MeV, and the ρ is close to saturating this bound.

In QCD with N flavors the masses of the vector mesons must decrease relative to f_π at least as fast as $1/\sqrt{N}$ as the number of flavors increases. This behavior is illustrated in one limit. Consider QCD with N flavors and N_C -colors in the limit $N_C \rightarrow \infty$ and $N \rightarrow \infty$, with N/N_C held fixed. In the fundamental theory at L loops, the largest contributions to any Green's function are proportional to $g^{2L} N_C^{L-k} N^k$ for $0 \leq k \leq L$ and where g is the strong coupling constant. As in conventional large- N_C QCD [19] this theory is well-behaved if we hold $g^2 N_C$ fixed. In this limit, the masses of resonances are fixed and f_π grows like $\sqrt{N_C}$ or, equivalently, $\sqrt{N}$. Therefore, the ratio of the masses of resonances to f_π falls like $1/\sqrt{N}$.

We wish to stress, however, that the behavior of Λ_χ with N is *not* dependent on taking the double limit. For example, when $N_C = 3$, we know that QCD is well defined for the six quarks that actually exist. If they were all light and f_π were fixed at 93 MeV, the masses of the analogs of the ρ would have to be less than 500 MeV!

In the one-family technicolor model, $N = 8$ and $f \approx 125$ GeV. The low energy prediction for the singlet spin zero channel would saturate unitarity at an energy of order 550 GeV. The arguments we have given imply that the resonances of this model must have masses of this order of magnitude. In technicolor phenomenology, however, one usually scales from QCD [23] by multiplying all mass scales by f_{TC}/f_π and applying large N_{TC} arguments [19]. When $N_{TC} = 3$ in the one-family model, this gives a technirho mass of about 1000 GeV. While we cannot say conclusively that there are *resonances* as light as 550 GeV, it is clear that the one-family model is quite different from QCD and that predictions based on scaling from QCD, e.g. [11], cannot be trusted in detail. In particular, if the masses of the resonances are much lighter than naively expected, then the estimates of “oblique” corrections from technicolor obtained by scaling from QCD [16], are untrustworthy.

Appendix A. Projection Operators and Identities for $SU(N)$

The following relations may be derived from the $SU(N)$ Fierz identities:

$$\begin{aligned}
f^{abe} f^{cde} &= d^{ace} d^{bde} - d^{ade} d^{bce} + \frac{2}{N}(\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc}) \\
f^{abe} d^{cde} &= d^{ade} f^{bce} + d^{ace} f^{bde} \\
d^{aef} d^{bef} &= \frac{N^2 - 4}{N} \delta^{ab} \\
d^{aef} d^{bfg} d^{cge} &= \frac{N^2 - 12}{2N} d^{abc} \\
d^{aef} d^{bfg} d^{cgh} d^{dhe} &= \frac{N^2 - 4}{N^2} (\delta^{ab} \delta^{cd} + \delta^{ad} \delta^{bc}) + \frac{N^2 - 16}{4N} (d^{abe} d^{cde} + d^{ade} d^{bce}) \\
&\quad - \frac{N}{4} d^{ace} d^{bde}
\end{aligned} \tag{A.1}$$

The operators that project a state $|\pi^a \pi^b\rangle$ onto the various irreducible representations of $SU(N)_V$ are

$$\begin{aligned}
P_\Delta &= \frac{1}{N^2 - 1} \delta^{ab} \delta^{cd} \\
P_F &= \frac{1}{N} (d^{ace} d^{bde} - d^{ade} d^{bce}) + \frac{2}{N^2} (\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc}) \\
P_D &= \frac{N}{N^2 - 4} d^{abe} d^{cde} \\
P_Y &= \frac{N - 2}{4N} (\delta^{ac} \delta^{bd} + \delta^{ad} \delta^{bc}) + \frac{N - 2}{2N(N - 1)} \delta^{ab} \delta^{cd} \\
&\quad - \frac{1}{4} (d^{ace} d^{bde} + d^{ade} d^{bce}) + \frac{N - 4}{4(N - 2)} d^{abe} d^{cde} \\
P_X &= \frac{N + 2}{4N} (\delta^{ac} \delta^{bd} + \delta^{ad} \delta^{bc}) - \frac{N + 2}{2N(N + 1)} \delta^{ab} \delta^{cd} \\
&\quad + \frac{1}{4} (d^{ace} d^{bde} + d^{ade} d^{bce}) - \frac{N + 4}{4(N + 2)} d^{abe} d^{cde} \\
P_T &= P_T^* = \frac{N^2 - 4}{4N^2} (\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc}) \\
&\quad - \frac{1}{2N} (d^{ace} d^{bde} - d^{ade} d^{bce}) + \frac{i}{4} (d^{ace} f^{bde} - d^{ade} f^{bce}) .
\end{aligned} \tag{A.2}$$

Using the identities (A.1), one can verify that these are projection operators satisfying

$$P_I^{a,b;c,d} P_K^{c,d;g,h} = \delta_{IK} P_I^{a,b;g,h}, \tag{A.3}$$

and

$$\Sigma_I P_I^{a,b;c,d} = \delta^{ac} \delta^{bd}. \tag{A.4}$$

We may write the amplitude in terms of the seven $SU(N)_V$ channels:

$$a(s, t, u)^{a,b;c,d} = \sum_I a_I(s, t, u) P_I^{a,b;c,d}, \quad (\text{A.5})$$

where I runs over $\Delta, F, D, Y, X, T, \bar{T}$, yielding eqn. (2.3).

Appendix B. Renormalization of $\mathcal{L}_4$

Calculating at one loop, there are renormalizations to the pion mass, the decay constant, and the pion wave function. We define

$$\begin{aligned} \pi^a &= Z_\pi^{-1/2} \pi_0^a \\ m &= Z_m m_0 \\ f &= Z_f f_0 \end{aligned} \quad (\text{B.1})$$

where the subscript 0 denotes the bare quantities. The calculation of the mass and wave-function renormalizations is straightforward. We demand that the pion propagator have a pole of residue 1 at m^2 . There is only one one-loop diagram which contributes. In addition some terms in $\mathcal{L}_4$ make a contribution, and the total is

$$\begin{aligned} Z_\pi &= 1 - \frac{m^2}{f^2} \left(8N\ell_4 + 8\ell_5 + \frac{N}{3} \left(D + \frac{1}{16\pi^2} - \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} \right) \right) \\ Z_m &= 1 - \frac{m^2}{f^2} \left(4N\ell_4 + 4\ell_5 - 8N\ell_6 - 8\ell_8 + \frac{1}{2N} \left(D + \frac{1}{16\pi^2} - \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} \right) \right) \end{aligned} \quad (\text{B.2})$$

Here μ is scale of dimensional regularization and D is the quantity $\frac{1}{16\pi^2}(2/\epsilon - \gamma_E + \log 4\pi)$.

To renormalize f , we must define the axial current. To do this we replace $\partial^\mu \Sigma$ by $(\partial^\mu - ia^\mu)\Sigma - i\Sigma a^\mu$ in $\mathcal{L}$. The axial current A^μ is what multiplies a^μ in $\mathcal{L}$; at lowest order it is $A_\mu^a = (if^2/2)\text{tr}(T^a(\Sigma^\dagger \partial_\mu \Sigma - \Sigma \partial_\mu \Sigma^\dagger))$. The decay constant f is then defined by

$$\langle 0 | A^{a\mu} | \pi^b \rangle = if p^\mu \delta^{ab} \quad (\text{B.3})$$

The operators multiplying $\ell_1 \dots \ell'_3$ make a contribution to A^μ , but not to the matrix element in (B.3). The only contributions in $\mathcal{L}_4$ are from the ℓ_4 and ℓ_5 terms. In addition there is a one-loop diagram from $\mathcal{L}_2$. The result is

$$Z_f = 1 + \frac{m^2}{f^2} \left(4N\ell_4 + 4\ell_5 + \frac{N}{2} \left(D + \frac{1}{16\pi^2} - \frac{1}{16\pi^2} \log \frac{m^2}{\mu^2} \right) \right) \quad (\text{B.4})$$

Finally, the infinities in the computation of the scattering amplitudes can all be absorbed by defining renormalized quantities $\ell(\mu)$ by

$$\begin{aligned}\ell_1(\mu) &= \ell_1 + \frac{1}{32}D & \ell_2(\mu) &= \ell_2 + \frac{1}{16}D & \ell_3(\mu) &= \ell_3 + \frac{N}{48}D \\ \ell'_3(\mu) &= \ell'_3 + \frac{N}{96}D & \ell_4(\mu) &= \ell_4 + \frac{1}{16}D & \ell_5(\mu) &= \ell_5 + \frac{N}{16}D \\ \ell_6(\mu) &= \ell_6 + \left(\frac{1}{32} + \frac{1}{16N^2}\right)D & \ell_8(\mu) &= \ell_8 + \left(\frac{N}{32} - \frac{1}{8N}\right)D\end{aligned}\tag{B.5}$$

This computation does not yield the renormalization of ℓ_7 because that term makes no contribution to $\pi\pi \rightarrow \pi\pi$ scattering. Gasser and Leutwyler have computed the finite parts of $A(s, t, u)$ for the case of $N = 2$ and our formula agrees with theirs³. The results in (B.5) agree with their computation for the case of $N = 3$, after one eliminates the operator corresponding to ℓ'_3 .

Acknowledgements.

We thank Andrew Cohen, Howard Georgi, Ken Lane, Stephen Selipsky, and Elizabeth Simmons for useful conversations and German Valencia for providing us with the data compiled in ref. [22]. R.S.C. acknowledges the support of a NSF Presidential Young Investigator Award and of an Alfred P. Sloan Foundation Fellowship. M.G. thanks the Texas National Research Laboratory Commission for support under a Superconducting Super Collider National Fellowship. This research is supported in part by the National Science Foundation under grants PHY-87-14654 and PHY-90-57173, by the Texas National Research Laboratory Commission, under grants RGFY9106 and RGFY91B6, and by the Department of Energy, under contract numbers DE-AC02-89ER40509 and DE-FG02-91ER40676.

³ They do not use the $\overline{MS}$ prescription. Their subtractions are proportional to $D + 1/16\pi^2$ instead of D .

References

- [1] S. Weinberg, Phys. Rev. **166** (1968) 1568;
S. Coleman, J. Wess, and B. Zumino, Phys. Rev. **177** (1969) 2239 ;
C. Callan, S. Coleman, J. Wess, and B. Zumino Phys. Rev. **177** (1969) 2246.
- [2] T. Appelquist and C. Bernard, Phys. Rev. D **22** (1980) 200 and Phys. Rev. D **23** (1981) 425;
A. Longhitano, Phys. Rev. D **22** (1980) 1166 and Nucl. Phys. **B231** (1984) 205.
- [3] M. S. Chanowitz and M. K. Gaillard, Nucl. Phys. **B261** (1985) 379.
- [4] M. S. Chanowitz, M. Golden, and H. Georgi, Phys. Rev. Lett. **57** (1986) 2344, and Phys. Rev. D **35** (1987) 1490.
- [5] S. Weinberg, Phys. Rev. D **19** (1979) 1277;
L. Susskind, Phys. Rev. D **20** (1979) 2619.
- [6] E. Eichten and K. Lane, Phys. Lett. **90B** (1980) 125;
S. Dimopoulos and L. Susskind, Nucl. Phys. **B155** (1979) 237.
- [7] E. Farhi and L. Susskind, Phys. Rev. D **20** (1979) 3404.
- [8] J. Bagger, S. Dawson, and G. Valencia, Phys. Rev. Lett. **67** (1991) 2256.
- [9] S. Weinberg, Physica **96A** (1979) 327;
see also H. Georgi and A. Manohar, Nucl. Phys. **B234** (1984) 189.
- [10] M. Soldate and R. Sundrum, Nucl. Phys. **B340** (1990) 1
- [11] K. Lane, in *Proceedings of the 1982 DPF Summer Study on Elementary Particle Physics and Future Facilities*, edited by R. Donaldson *et al*, (Fermilab, Batavia, Ill.), p. 222;
E. Eichten, I. Hinchliffe, G. Lane, and C. Quigg, Rev. Mod. Phys. **56** (1984) 579.
- [12] R. S. Chivukula and M. Golden, Phys. Lett. **267B** (1991) 233;
R. S. Chivukula, M. Golden, and M. V. Ramana, Phys. Rev. Lett. **68** (1992) 2883.
- [13] B. Lee, C. Quigg, and H. Thacker, Phys. Rev. Lett. **38** (1977) 883.
- [14] R. N. Cahn and M. Suzuki, Phys. Rev. Lett. **67** (1991) 169.
- [15] K. Jung and R. Willey, Phys. Rev. D **9** (1974) 3132;
A. Dobado and M. Herrero, Phys. Lett. **228B** (1989) 495;
A. Dobado, M. Herrero, and T. Truong, Phys. Lett. **235B** (1990) 129 and 134;
A. Dobado, Phys. Lett. **237B** (1990) 457.
- [16] M. Peskin, T. Takeuchi, Phys. Rev. Lett. **65** (1990) 964, and SLAC preprint SLAC-PUB-5618 (Nov 1991);
R. N. Cahn and M. Suzuki, Phys. Rev. D **44** (1991) 3641.
- [17] S. Coleman, *Aspects of Symmetry*, (Cambridge University Press, 1985) p21
- [18] Gasser and Leutwyler, Ann. Phys. (NY) **158** (1984) 142 and Nucl. Phys. **B250** (1985) 465.
- [19] G. 't Hooft, Nucl. Phys. **B72** (1974) 461.

- [20] L.-F. Li and H. Pagels, Phys. Rev. Lett. **26** (1971) 1089;
see also H. Pagels, Phys. Repts. **16C** (1975) 221.
- [21] See, for example, Shang-Keng Ma, *Modern Theory of Critical Phenomena* (Benjamin/Cummings, 1976)
- [22] J. F. Donoghue, C. Ramirez, and G. Valencia, Phys. Rev. D **38** (1988) 2195.
- [23] S. Dimopoulos, S. Raby, and G. Kane, Nucl. Phys. **B182** (1981) 77
- [24] N. Cason, *et al*, Phys. Rev. D **28** (1983) 1586;
V. Srinivasan, *et al*, Phys. Rev. D **12** (1975) 681;
L. Rosselet, *et al*, Phys. Rev. D **15** (1977) 574;
A. Belkov, *et al*, JETP Lett. 29 (1979) 597;
B. Hyams, *et al*, Nucl. Phys. **B73** (1974) 202;
E. Alekseeva, *et al*, Sov. Phys. JETP 55 (1982) 591 and JETP Lett. 29 (1979) 100 ;
W. Hoogland, *et al*, Nucl. Phys. **B69** (1974) 266;
J. Baton, *et al*, Phys. Lett. **33B** (1970) 525 and 528;
J. Prukop, *et al*, Phys. Rev. D **10** (1974) 2055;
D. Cohen, *et al*, Phys. Rev. D **7** (1973) 662.

Figure Captions

- Fig. 1. Data and lowest-order prediction for $Re\ a_{\Delta 0}$. The data is from a compilation in ref. [22] of the experimental results in ref. [24].
- Fig. 2. Data and lowest-order prediction as in fig. 1, for $|a_{F1}|$.
- Fig. 3. Data and lowest-order prediction as in fig. 1, $Re\ a_{X0}$.